



Product Information

TRIS Acetate-EDTA buffer

Product Number **T 9650**
Store at Room Temperature

Product Description

Synonym: TAE buffer

This product is a 10x concentrate containing 0.4 M Tris acetate (pH approximately 8.3) and 0.01 M EDTA. It is prepared with 18 megohm water and is 0.2 μ m filtered.

This product is suitable for agarose gel electrophoresis after dilution to the working concentration. This product has been analyzed for the absence of nucleases.

Tris-Acetate-EDTA (TAE) running buffer is a commonly used buffer for DNA agarose gel electrophoresis, and is especially useful in preparative work.¹ Compared to Tris-Borate-EDTA (TBE) and Tris-Phosphate-EDTA (TPE) buffers, double-stranded DNA tends to run faster in TAE. However, because TAE has the lowest buffering capacity of the three buffers, the buffering capacity can become exhausted during extended electrophoresis. Buffer circulation or replacement can remedy this situation.

The 1x TAE buffer is used both in the agarose gel and as a running buffer. Applied voltages of < 5 V/cm (the distance between the electrodes of the unit) are recommended for maximum resolution.² TAE buffer has been utilized in agarose gel electrophoresis of RNA.^{3,4} A study of free DNA solution mobility in TAE at various buffer concentrations, in the presence and absence of added NaCl, has been reported.⁵ The use of TAE buffer in a denaturing gradient gel electrophoresis method for broad-range mutation analysis has been described.⁶

Precautions and Disclaimer

For Laboratory Use Only. Not for drug, household or other uses.

References

1. Ogden, R. C., and Adams, D. A., Electrophoresis in agarose and acrylamide gels. *Methods Enzymol.*, **152**, 61-87 (1987).
2. Molecular Cloning: A Laboratory Manual, 3rd ed., Sambrook, J. and Russell, D. W., CSHL Press (Cold Spring Harbor, NY: 2001), pp. 5.8, 5.76, A1.16.
3. Loening, U. E., The fractionation of high-molecular-weight ribonucleic acid by polyacrylamide-gel electrophoresis. *Biochem. J.*, **102**, 251-257 (1967).
4. Masters, D. B., et al., High sensitivity quantification of RNA from gels and autoradiograms with affordable optical scanning. *Biotechniques*, **12(6)**, 902-906, 908-911 (1992).
5. Stellwagen, E., and Stellwagen, N. C., The free solution mobility of DNA in Tris-acetate-EDTA buffers of different concentrations, with and without added NaCl. *Electrophoresis*, **23(12)**, 1935-1941 (2002).
6. Hayes, V. M., et al., Improvements in gel composition and electrophoretic conditions for broad-range mutation analysis by denaturing gradient gel electrophoresis. *Nucleic Acids Res.*, **27(20)**, e29 (1999).

GCY/RXR 8/03

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.