



3050 Spruce Street
Saint Louis, Missouri 63103 USA
Telephone 800-325-5832 • (314) 771-5765
Fax (314) 286-7828
email: techserv@sial.com
sigma-aldrich.com

Product Information

Anti-phospho-Tau [pSer⁴²²]

Developed in Rabbit, Affinity Isolated Antibody

Product Number **T 7944**

Product Description

Anti-phospho-Tau [pSer⁴²²] is developed in rabbit using a synthetic phosphopeptide derived from the region of human tau that contains serine 422 as immunogen. The serum is affinity purified using epitope-specific affinity chromatography. The antibody is preadsorbed to remove any reactivity towards a non-phosphorylated tau.

Anti-phospho-Tau [pSer⁴²²] recognizes human, mouse and rat tau [pSer422] (45-68 kDa). It has been used in immunoblotting applications.

Tau is a microtubule-associated phosphoprotein (MAP), localized in neuronal axons. It promotes tubulin polymerization and stabilizes microtubules.¹ The biological activity of tau is regulated by its degree of phosphorylation.^{1,2} Hyperphosphorylated tau is the major protein of the paired helical filaments (PHFs), which make up the pathological neurofibrillary tangles of Alzheimer's disease (AD). The PHFs are also found in the lesions of other central nervous system disorders.^{3,4}

Tau phosphorylation involves numerous kinases: glycogen synthase kinase 3 β (GSK-3 β), MARK kinase, MAP kinase, protein kinase A and C, cyclin-dependent kinase 5 (Cdk5), p38 kinase, c-Jun N-terminal kinase, and casein kinase II.^{1,2,5,6,7} Combined tau protein kinase II (TPKII), which consists of Cdk5 and GSK-3 β , is the most potent phosphorylation agent indirectly involved in the regulation of the phosphorylation state of tau in neuronal cells.^{6,8} In addition, tau is phosphorylated *in vitro* by osmotic cellular stress, which activates the stress-activated protein kinases (SAPKs).

To date, a total of 25 abnormal phosphorylation sites have been identified on hyperphosphorylated tau in AD brain.¹⁰ Normal tau has approximately eight phosphorylation sites. The abnormal phosphorylation occurs usually on serine and threonine residues. Specifically, TPKII phosphorylates serines 202 and 404. GSK-3 β transfection phosphorylates serines 199, 202, 235, 396, 404 and 413, and threonines 205 and 231.

These sites are among the major abnormal phosphorylation sites of tau.¹¹ Phosphorylation on these sites reduces the ability of a given tau species to promote microtubule self-assembly.^{11,12} Okadaic acid increases phosphorylation at threonine 231 and serines 235, 396 and 404. Phosphorylated serine 422 was found in the biopsies of brains from patients with Down syndrome, amyotrophic lateral sclerosis, corticobasal degeneration, and Pick's disease. It was absent from a control group of normal brains.¹³

The opposite process, tau dephosphorylation, is controlled by different protein phosphatases expressed in neurons. Protein phosphatases PP2A and PP2B efficiently dephosphorylate tau *in vitro* and restore biological activity in the assembly of microtubules.^{3,10,14}

Recently it was discovered that propyl isomerase (Pin 1) interacts with tau hyperphosphorylated on threonine 231 and restores the ability of tau to bind to microtubules.

Reagent

The antibody is supplied as a solution in Dulbecco's phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.3, with 50% glycerol, 1.0 mg/ml BSA (IgG and protease free), and 0.05% sodium azide.

Precautions and Disclaimer

Due to the sodium azide content a material safety data sheet (MSDS) for this product has been sent to the attention of the safety officer of your institution. Consult the MSDS for information regarding hazardous and safe handling practices.

Storage/Stability

Store at -20 °C. For extended storage, upon initial thawing, freeze in working aliquots. Avoid repeated freezing and thawing to prevent denaturing the antibody. Working dilution samples should be discarded if not used within 12 hours.

Product Profile


The supplied reagent is sufficient for 10 immunoblots.

A recommended working dilution of 1:1000 is determined by immunoblotting.

Note: In order to obtain best results in different techniques and preparations we recommend determining optimal working concentration by titration test.

Peptide Competition

- Human recombinant Tau stimulated with GSK-3 β (1 μ g per μ g Tau) for 45 minutes was added to background extracts, resolved by SDS-PAGE on a 10% Tris-glycine gel and transferred to PVDF.
- Membranes were blocked with a 5% BSA-TBST buffer for 1 hour at room temperature.
- After blocking, membranes were left untreated (lanes 1-4) or treated with lambda (λ) phosphatase (lane 5), and preincubated with different peptides as follow:
 Lane 1, 5 no peptide
 Lane 2 non phosphorylated peptide corresponding to the immunogen
 Lane 3 a generic phosphoserine containing peptide
 Lane 4 immunogen
- After preincubation membranes were incubated with Anti-phospho-Tau [pSer⁴²²] in a 3% BSA-TBST buffer for two hours at room temperature.
- After washing, membranes were incubated with goat F(ab')₂ anti-rabbit IgG alkaline phosphatase and signals were detected using the Pierce SuperSignal™ method.



The data show that only the peptide corresponding to Tau [pSer⁴²²] blocks the antibody signal, thereby demonstrating the specificity of the antibody. The data also show that phosphatase stripping eliminates the signal, verifying that the antibody is phospho-specific.

References

- Haque, N., et al., Regulation of expression, phosphorylation and biological activity of tau during differentiation in SY5Y cells., *Brain Res.*, **838**, 69-77 (1999).
- Jenkins, S.M., et al., Modulation of tau phosphorylation and intracellular localization by cellular stress., *Biochem. J.*, **345**, 263-270 (2000).
- Gong, C.X., et al., Phosphorylation of microtubule-associated protein tau is regulated by protein phosphatase 2A in mammalian brain. Implications for neurofibrillary degeneration in Alzheimer's disease., *J. Biol. Chem.*, **275**, 5535-5544 (2000).
- Goedert, M., Alzheimer's disease. Pinning down phosphorylated tau., *Nature*, **399**, 739-740 (1999).
- Maccioni, R.B., et al., The protein kinase Cdk5. Structural aspects, roles in neurogenesis and involvement in Alzheimer's pathology., *Eur. J. Biochem.*, **268**, 1518-1527(2001).
- Arioka, M. et al., Tau protein kinase II is involved in the regulation of the normal phosphorylation state of tau protein., *J. Neurochem.*, **60**, 461-468 (1993).

7. Anderton, B.H., et al., Sites of phosphorylation in tau and factors affecting their regulation., *Biochem. Soc. Symp.*, **67**, 73-80 (2001).
8. Michel, G., et al., Characterization of tau phosphorylation in glycogen synthase kinase-3 β and cyclin dependent kinase-5 activator (p23) transfected cells., *Biochem. Biophys. Acta*, **1380**, 177-182 (1998).
9. Lund, E.T., et al., Characterization of the *in vitro* phosphorylation of human tau by tau protein kinase II (cdk5/p20) using mass spectrometry., *J. Neurochem.*, **76**, 1221-1232 (2001).
10. Gong, C.X., et al., Alzheimer's disease abnormally phosphorylated tau is dephosphorylated by protein phosphatase-2B (calcineurin)., *J. Neurochem.*, **62**, 803-806 (1994).
11. Takashima, A., et al., Activation of tau protein kinase I/glycogen synthase kinase-3 β by amyloid β peptide (25-35) enhances phosphorylation of tau in hippocampal neurons., *Neurosci. Res.*, **31**, 317-323 (1998).
12. Utton, M.A., et al., Phosphorylation of tau by glycogen synthase kinase 3 β affects the ability of tau to promote microtubule self-assembly., *Biochem. J.*, **323**, 741-747 (1997).
13. Bussiere, T., et al., Phosphorylated serine422 on tau proteins is a pathological epitope found in several diseases with neurofibrillary degeneration., *Acta Neuropathol. (Berlin)*, **97**, 221-230 (1999).
14. Wang, J.Z., et al., Restoration of biological activity of Alzheimer abnormally phosphorylated tau by dephosphorylation with protein phosphatase-2A, -2B and -1., *Brain. Res. Mol. Brain Res.*, **38**, 200-208 (1996).

SuperSignal is a registered trademark of Pierce Biotechnology, Inc.

AH,PHC 03/05-1

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.