

Product Information

Monoclonal Anti-c-Myc antibody produced in mouse

Clone 9E10, purified immunoglobulin, buffered aqueous solution

M4439

Product Description

The human *c-myc* proto-oncogene is the human cellular homologue of the avian *v-myc* gene that is present in several leukemogenic retroviruses.^{1,2} Increased *c-myc* expression has been described in various human tumors that occur by mechanisms such as gene amplification and chromosomal translocation.³ The gene encodes a polypeptide with a predicted molecular weight of 49 kDa, which shows aberrant electrophoretic mobility on polyacrylamide gel electrophoresis to give an apparent molecular weight of around 62 kDa (p62c-myc).⁴ p62c-myc is associated mainly with the cell nucleus, where it exerts its normal and oncogenic functions.^{1,3}

An epitope located within amino acids 410-419, which contains the sequence EQKLISEEDL of human c-Myc, has been widely used as a tag in many expression vectors, to enable the expression of proteins as c-Myc-tagged fusion proteins.⁵ Epitope tags provide a method to localize gene products in a variety of cell types, to study the topology of proteins and protein complexes, and to identify associated proteins. These tags also allow characterization of newly identified, low abundance or poorly immunogenic proteins when protein-specific antibodies are not available.^{5,6}

Monoclonal Anti-c-Myc (mouse IgG1 isotype) is derived from the 9E10 hybridoma, which is produced by fusion of mouse myeloma cells and splenocytes from BALB/c mice. A synthetic peptide corresponding to residues 408-439 of the human p62c-myc protein conjugated to KLH was used as immunogen.¹ The isotype was determined using the Roche IsoStrip™ Mouse Monoclonal Antibody Isotyping Kit (Cat. No. 11493027001).

Monoclonal Anti-c-Myc recognizes an epitope located within the sequence EQKLISEEDL (residues 410-419) of the product of the human oncogene *c-myc*.⁷ The antibody recognizes the c-Myc tag sequence when it is expressed at either the amino or the carboxyl terminus of the fusion protein. The antibody reacts specifically with c-Myc-tagged fusion proteins in immunoblotting, immunoprecipitation and immunofluorescence applications. Reaction of the antibody in immunoblotting is inhibited by the c-Myc peptide (Cat. No. M2435).⁶

The antibody reacts with both components of the p62c-myc-p64c-myc doublet, by immunoblotting.^{1,7} The antibody is also useful in immunohistochemical labeling of the c-Myc oncoprotein in formalin-fixed paraffin-embedded tissue sections, applying light microscopy⁸ and electron microscopy.^{4,8} Additional applications of the product include ELISA.¹ The antibody cross-reacts with human p62/64^{c-myc}, but fails to recognize the chicken p11^{gag-myc} protein present in MC29 virus-transfected quail fibroblasts, nor does it react with the mouse p64/66^{c-myc} protein.¹ Nevertheless, weak reaction with murine c-myc may be seen when the antibody is used at high concentration.

Several theses⁹⁻¹⁴ and dissertations¹⁵⁻³⁴ cite use of this M4439 product in their research protocols.

Reagent

Monoclonal Anti-c-Myc is supplied as purified IgG in 0.01 M phosphate buffered saline (pH 7.4), containing 15 mM sodium azide as a preservative.

Antibody concentration: ~ 2 mg/mL (exact information on lot-specific Certificate of Analysis).

Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For extended storage, freeze in working aliquots at -20 °C. Repeated freezing and thawing is not recommended. Storage in a frost-free freezer is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilutions should be discarded if not used within 12 hours.

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazardous and safe handling practices.

Product Profile

A minimum working dilution of 1:5000 is determined by immunoblotting of an *E. coli* extract expressing a recombinant c-Myc-tagged fusion protein. The tagged protein was detected using a chemiluminescent substrate.

Note: To obtain optimal results using different techniques and preparations, we recommend determining the optimal working dilution by titration.

Immunoprecipitation

1-2 µg of the antibody can immunoprecipitate a c-Myc fusion protein from transfected mammalian cell lysates or bacterial extracts.

Note: To obtain best results and assay sensitivity in different techniques and preparations, we recommend determining optimal working dilutions by titration.

Procedure

Procedure for Immunoblotting

Note: All incubation steps should be performed at room temperature.

1. Separate c-Myc-tagged proteins from sample lysates using a standard sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) protocol. Load 2.5-20 µg of total lysate protein per lane. The amount of lysate to be loaded per lane depends on the level of protein expression. Thus, the optimum loading may vary between preparations.
2. Transfer proteins from the gel to a nitrocellulose membrane.
3. Block the membrane using a solution of 5% non-fat dry milk in phosphate buffered saline (PBS, Cat. No. D8537) for at least 60 minutes.

4. Wash the membrane three times for 5 minutes each in PBS containing 0.05% Tween® 20 (PBS-T, Cat. No. P3563).
5. Incubate the membrane with Monoclonal Anti-c-Myc as the primary antibody for 60-120 minutes, using an optimized concentration in PBS-T.
6. Wash the membrane three times for 5 minutes each in PBS-T.
7. Incubate the membrane with anti-mouse IgG Peroxidase conjugate (Cat. Nos. A9917, A3682, or A2304) or with anti-mouse Fab Alkaline Phosphatase conjugate (Cat. Nos. A1293, A2179 or A1682) as the secondary antibody, at the recommended concentration in PBS-T, for 60 minutes. Adjust the antibody concentration to maximize detection sensitivity and to minimize background.
8. Wash the membrane three times for 5 minutes each in PBS-T.
9. Treat the membrane with a peroxidase or an alkaline-phosphatase substrate.

Procedure for Indirect Immunofluorescent Staining of Cultured Cells

Note: All incubation steps should be performed at room temperature, except Steps 1 and 3.

1. Grow transfected cultured cells which express a c-Myc-tagged protein on sterile coverslips at 37 °C.
2. Wash the cells briefly in PBS.
3. Fix the cells:
 - a. First with -20 °C methanol (10 minutes),
 - b. Then with -20 °C acetone (1 minute).
4. Wash coverslips twice in PBS (5 minutes each wash).
5. Incubate the coverslips cell-side-up with Monoclonal Anti-c-Myc in PBS containing 1% BSA, for 60 minutes.
6. Wash three times in PBS (5 minutes each wash).
7. Incubate the coverslips, cell-side-up, with anti-mouse Fab, FITC conjugate (Cat. No. F4018 or F8771) as the secondary antibody, at the recommended dilution, in PBS containing 1% BSA, for 30 minutes.
8. Wash three times in PBS (5 minutes each wash).
9. Add one drop of aqueous mounting medium on the coverslip and invert carefully on a glass slide. Avoid air bubbles.
10. Examine using a fluorescence microscope with appropriate filters.

Note: Blocking with PBS containing 1% BSA for 10 minutes at room temperature, followed by draining prior to Step 5, may minimize non-specific adsorption of the antibody.

Procedure for Immunoprecipitation

1. Centrifuge 40 μ L of Protein G-Sepharose® beads, 1:1 suspension (Cat. No. P3296) for 1 minute at 12,000 \times g. Then wash twice with 1 mL RIPA buffer (50 mM Tris base, pH 7.4, containing 0.25% (w/v) deoxycholate, 1% Igepal® CA-630, 150 mM NaCl, and 1 mM EDTA) at 4 °C.
2. Add Monoclonal Anti c-Myc diluted in PBS. Incubate with continual inversion for 60 minutes.
3. Centrifuge for 1 minute at 12,000 \times g. Then wash twice with 1 mL RIPA buffer at 4 °C.
4. Add 0.1-1.0 mL of cell extract containing the c-Myc-tagged protein to the beads (see **Note**). Incubate from 2 hours to overnight at 4 °C, with continual inversion.

Note: The amount of cell extract required depends on the level of expression of the tagged protein and the specific application.

5. Separate the beads by centrifugation and remove the supernatant.
6. Wash the beads five times with 1 mL PBS each. Vortex (mix) the beads in the PBS. Then separate the beads by brief centrifugation.
7. Resuspend the pellet in 25 μ L 2 \times SDS-PAGE sample buffer. Boil the sample for 5 minutes and centrifuge. The sample is ready to be loaded on a SDS-PAGE gel.

References

1. Evan, G. *et al.*, *Mol. Cell Biol.*, **5(12)**, 3610-3616 (1985).
2. Alitalo, K. *et al.*, *Nature*, **306(5940)**, 274-277 (1983).
3. Pelengaris, S. *et al.*, *Curr. Opin. Genet. Dev.*, **10(1)**, 100-105 (2000).
4. Robertson, D. *et al.*, *J. Histochem. Cytochem.*, **43(5)**, 471-480 (1995).
5. Olins, P.O., and Lee, S.C., *Curr. Opin. Biotechnol.*, **4(5)**, 520-525 (1993).
6. Jarvik, W., and Telmer, C.A., *Annu. Rev. Genet.*, **32**, 601-618 (1998).
7. Campbell, A. *et al.*, *J. Biol. Chem.*, **267(13)**, 9321-9325 (1992).
8. Royds, J.A. *et al.*, *J. Pathol.*, **166(3)**, 225-233 (1992).
9. Yang, Dian, "Biochemistry of a mutant, cancer-causing DNA licensing protein MCM4". Cornell University, B.S. thesis (honors), p. 11 (May 2011).
9. Taylor, Rebecca J., "Understanding Collagen-I Folding and Misfolding". Massachusetts Institute of Technology, B.S. thesis, p. 30 (June 2015).
10. Welte, Emily, "Keratins in Skeletal Muscle: Effects of Altering Expression of Type I/Type II Keratin Pairs". University of Maryland, Baltimore, M.S. thesis, p. 70 (2014).
11. Garcia, Sara, "Investigating The Role Of Parp-1 In HIV-1 Replication". University of Texas at El Paso, M.S. thesis, p. 21 (May 2018).
12. Choi, Myoung Soo, "Identification of CALML4 as a Novel Component of the Intermicrovillar Adhesion Complex that Regulates Intestinal Brush Border Assembly". University of Toledo, M.S. thesis, p. 22 (August 2018).
13. Saado, Indira, "Functional Elucidation of Pathogen Triggered Immunity (PTI) Inhibiting Effectors of *Ustilago maydis*". Universität Wien (University of Vienna), M.Sc. thesis, p. 53 (2018).
14. Horwich, Michael D., "Small RNA sorting in *Drosophila* produces chemically distinct functional RNA-protein complexes". University of Massachusetts Worcester, Ph.D. dissertation, p. 70 (2008).
15. Liu, Fen, "Misfolded cytosolic proteins are trafficked through the endoplasmic reticulum for degradation: the molecular and cellular mechanisms for the turnover of human NAT1 R64W and parkin R42P". University of Minnesota, Ph.D. dissertation, p. 101 (April 2009).
16. Ramsey, Chenere P., "Studies of DJ-1, Parkin and Alpha-Synuclein Give Insights into Plausible Mechanisms for Parkinson's Disease Pathogenesis". University of Pennsylvania, Ph.D. dissertation, p. 154 (2010).
17. Samel, Anke, "Identification and characterization of posttranslational modifications on CenH3 in *Saccharomyces cerevisiae*". Universität Duisburg-Essen, Dr. rer. nat. dissertation, p. 45 (February 2012).
18. Gegg, Moritz A., "Identification and characterization of the novel planar cell polarity gene *Flattop (Fltp)*". Technische Universität München, Ph.D. dissertation, p. 113 (2012).
19. Shveygert, Mayya, "Regulation of Eukaryotic Translation Initiation by Signal Transduction". Duke University, Ph.D. dissertation, p. 68 (2012).
20. Waters, Aoife Mary, "Genetic and functional dissection of ciliary genes". University College London, Ph.D. dissertation, p. 102 (2012).

21. Reinhard, Christian, "HIV-1 innate immune detection and evasion". University of Geneva, Ph.D. dissertation, p. 105 (2013).
22. Carmona-Mora, Paulina, "The molecular role of GTF2IRD1: a protein involved in the neurodevelopmental abnormalities of Williams-Beuren syndrome". University of New South Wales, Ph.D. dissertation, p. 49 (May 2015).
23. Rubel, Carrie E., "Defining the Molecular Mechanisms of Ubiquitin Proteasome System Dysfunction as a Driver of Disease: CHIP mutation in SCAR16". University of North Carolina Chapel Hill, Ph.D. dissertation, p. 183 (2015).
24. Zhang, Jin, "The subproteome of mitoBK_{Ca} channels from cardiomyocytes reveals insights into its mitochondrial import mechanism and function". University of California Los Angeles, Ph.D. dissertation, p. 29 (2016).
25. Xu, Xiaosu, "Conserved Modulation of the Constitutive Photomorphogenic E3 Ubiquitin Ligase Activity by the bHLH Transcription Factors, PHYTOCHROME INTERACTING FACTORS". University of Texas at Austin, Ph.D. dissertation, p. 30 (May 2016).
26. Aslam, Kiran, "Deciphering the role of Hsp31 as a multitasking chaperone". Purdue University, Ph.D. dissertation, p. 141 (August 2016).
27. Black, Hannah Lucy, "Impact of tyrosine phosphorylation of Syntaxin 4 and Munc18c on GLUT4 translocation". University of Glasgow, Ph.D. dissertation, p. 62 (September 2016).
28. Urakova, Nadezda, "Identifying molecular virulence factors of rabbit haemorrhagic disease virus". University of Canberra, Ph.D. dissertation, pp. 13, 35, 154, 175 (January 2017).
29. Withers, Catherine Nicole Kaminski, "Rad GTPase: identification of novel regulatory mechanisms and a new function in modulation of bone density and marrow adiposity". University of Kentucky, Ph.D. dissertation, p. 65 (2017).
30. Lin, Feng, "Molecular functions of cell plate-associated phosphoinositides during plant somatic cytokinesis". Martin-Luther-Universität Halle-Wittenberg, Dr. rer. nat. dissertation, p. 127 (September 2018).
31. Le, Uyen Quynh Nguyen, "Defining the role of tumor suppressor Dear1 in the acquisition of mammary stem/progenitor cell properties". University of Texas M.D. Anderson Cancer Center, p. 107 (December 2018).
32. Yan, Jiong, "Hepatic xenobiotic receptors in the ubiquitin-proteasome system". University of Pittsburgh, Ph.D. dissertation, pp. 29, 104 (2018).
33. Yang, Qian, "The Roles of Codon Usage in Translational and Transcriptional Regulation on Gene Expression". University of Texas Southwestern Medical Center at Dallas, Ph.D. dissertation, p. 31 (December 2020).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at [SigmaAldrich.com/techservice](https://www.sigmaaldrich.com/techservice).

Standard Warranty

The applicable warranty for the products listed in this publication may be found at [SigmaAldrich.com/terms](https://www.sigmaaldrich.com/terms).

Contact Information

For the location of the office nearest you, go to [SigmaAldrich.com/offices](https://www.sigmaaldrich.com/offices).

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the U.S. and Canada.

MilliporeSigma, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2021 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

M4439dat Rev 11/21 NV,GCY